

TABLE 4

PREDICTORS OF INTIMATE PARTNER ABUSE COMMUNICATION  
BETWEEN ABUSED WOMEN PATIENTS AND THEIR MEDICAL  
CLINICIANS

Factor	N	Ever Communicated N (%)	Adjusted Odds Ratio (95% CI)
<b>Age</b>			
18-29	104	37 (36)	1.0
30-39	159	66 (42)	1.34 (0.86, 2.11)
40-46	112	56 (50)	1.13 (0.77, 1.65)
<b>Ethnicity</b>			
Non-Latina white	137	56 (41)	1.0
African American	125	59 (47)	1.77 (1.08, 2.92)*
Latina	108	41 (38)	0.88 (0.53, 1.47)
<b>Birthplace</b>			
United States	276	127 (46)	1.0
Other	99	32 (32)	0.57 (0.33, 0.99)*
<b>Education, years</b>			
<12	114	42 (37)	1.0
12	116	49 (42)	1.40 (0.89, 2.19)
>12	142	66 (46)	(0.66, 1.51)
<b>Regular clinician</b>			
No	144	47 (33)	1.0
Yes	226	112 (50)	1.16 (.86, 1.57)
<b>Direct clinician questioning</b>			
No	261	66 (25)	1.0
Yes	104	88 (85)	4.59 (3.22, 6.55)*
<b>Perception that clinicians have no time</b>			
	178	63 (35)	0.91 (0.66, 1.25)
<b>Perception that clinicians have no interest</b>			
	123	40 (33)	0.96 (0.69, 1.33)
<b>Concerns about confidentiality</b>			
	114	34 (30)	0.68 (0.48, 0.94)*

NOTE: As reported by study participants in San Francisco, 1997  
CI denotes confidence interval.